



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

There were several of these birds on the grounds, and their singing was so much in evidence that boys took it up and could often be heard whistling an imitation.

L. B. CUSHMAN.

North East, Pa., Dec. 30, 1913.

Sandpiper Notes.

The fall migration of 1913 proved to be rather unusual in regard to the sandpiper movements. Many species lingered later than usual in this locality and several new species were noted. Immense mixed flocks of shore birds fed on the mud flats around the bayous from Aug. 20 to Sept. 12. In these flocks the usual Pectoral, Semipalmated, Least, and Solitary Sandpipers and Lesser Yellowlegs were abundant. In addition Baird's Sandpipers (*Pisobia bairdi*) were present in considerable numbers from Aug. 22-27, and several specimens were taken. One Greater Yellowlegs (*Totanus melanoleucus*) was noted Oct. 6. From the 12th of September the numbers decreased rapidly, although a few individuals remained much later. The last records for the more common species are as follows: Pectoral Sandpiper (*Pisobia maculata*), Nov. 12; White-rumped Sandpiper (*Pisobia fuscicollis*), Nov. 7; Least Sandpiper (*Pisobia minutilla*), Oct. 12; Semipalmated Sandpiper (*Ereunetes pusillus*), Oct. 12; Yellow-legs (*Totanus flavipes*), Nov. 1; Solitary Sandpiper (*Hedromus solitarius solitarius*), Oct. 5; and Spotted Sandpiper (*Actitis macularia*), Oct. 8.

On the 6th of October, after the bulk of the shore birds had departed, a flock of four Red-backed Sandpipers (*Pelidna alpina sakhalina*) were noted feeding on the deserted mud flats and one was taken. The birds were tame and unsuspicious and allowed a close approach. On the 10th the ponds were visited again and a flock of about sixty found in the same place. Four were taken at this time. All authorities to which I have access state that this species is uncommon in the interior in the fall and it was a surprise to me to find them so numerous at this time. This is the first time that I have noted this form in the fall. A few lingered until the 25th of October, when the last one was seen.

IRA N. GABRIELSON, Marshalltown, Iowa.

"The Guide to Nature."

Many magazines, hundreds of schools and thousands of teachers and parents have tried to instruct children in a knowledge of nature. Yet the really natural child takes to nature for enjoyment like a duck to water.

Why urge the duck, why compel it to go into the water? When we destroy spontaneity and liberty, we prevent enjoyment and all consequent benefit. "We love the things that love us."

It is, however, not nature nor even natural science as a matter of instruction, as the adult understands it, that the child wants, but the fun of seeing things. Where is the boy or girl that is not pleased by the